


Isothermal Amplification

AMPLIFICATION AT A SINGLE TEMPERATURE



What is isothermal amplification?

The Polymerase Chain Reaction (PCR) is a well-known approach for amplifying a specific DNA or RNA (RT-PCR) sequence. PCR involves the reiterative cycling of a reaction cocktail between different temperatures to achieve amplification. As routine as PCR is in the molecular biology and molecular diagnostics laboratory, there are other methods of sequence-specific nucleic acid amplification.

These alternative approaches often do not require changing the reaction temperature and are referred to as isothermal amplification methods. Isothermal amplification methods are varied and have different advantages. In general, isothermal techniques are extremely fast and do not require thermocyclers, making them particularly well suited for field applications and point-of-care molecular diagnostics assays.

Advantages

- Fast
- Minimal equipment required
- Robust reactions in the presence of inhibitors
- Options for simplified optical or quantitative detection








Did you know that many of these products can be purchased in large volumes and custom formats? Learn more at www.neb.com/customizedsolutions

Not sure which product will work best for your experiment?

NEB has engineered improved versions of *Bst* DNA Polymerase-based products for isothermal amplification. Use this chart to determine which product will work best for your needs.

Comparison of *Bst* DNA Polymerases

	<i>Bst</i> -XT WarmStart NEB #M9204	<i>Bst</i> 2.0 WarmStart® NEB #M0538	<i>Bst</i> 3.0 NEB #M0374
 Amplification speed	★★★★★	★★★	★★★★★
 Specificity	★★★★★	★★★★★	★★
 Room temperature set-up?	Enabled	Enabled	Not recommended
 Optimal LAMP temperature	50–70°C	60–70°C	55–72°C
 Available glycerol-free?	Yes NEB #M9205	Yes NEB #M0402	Yes NEB #M0443

- ★★★★★ Optimal, recommended product for selected application
 ★★ or ★★★ Will perform for selected application
 ★ May perform but is not recommended



View our **NEB TV webinar** to learn about the benefits of *Bst*-XT WarmStart DNA Polymerase.

Loop-mediated Isothermal Amplification (LAMP)

LAMP uses 4-6 primers recognizing 6-8 distinct regions of target DNA for a highly specific amplification reaction. A strand-displacing *Bst* DNA polymerase initiates synthesis and 2 specially designed primers form “loop” structures to facilitate subsequent rounds of amplification through extension on the loops and additional annealing of primers. DNA products are very long (>20 kb) and formed from numerous repeats of the short (80–250 bp) target sequence, connected with single-stranded loop regions in long concatamers. These products are not typically appropriate for downstream manipulation, but target amplification is so extensive that numerous modes of detection are possible. Real-time fluorescence detection using intercalating dyes or probes, lateral flow and visual detection are all directly compatible with monitoring a LAMP reaction. Instrumentation for LAMP typically requires consistent heating to the desired reaction temperature and, where needed, real-time fluorescence for quantitative measurements.

RECOMMENDED PRODUCTS

WarmStart® Fluorescent LAMP/RT-LAMP Kit (with UDG)
(NEB #E1708)


WarmStart Colorimetric LAMP 2X Master Mix with UDG
(NEB #M1804)

LyoPrime WarmStart™ Fluorescent LAMP/RT-LAMP Mix (with UDG)
(NEB #L4401)

WarmStart RTx Reverse Transcriptase
(NEB #M0380)

***Bst* DNA Polymerase**
(NEB #M0275, M0328, M0374, M0537, M0538, M9204)

Overview of Loop-Mediated Isothermal Amplification (LAMP)



Whole Genome Amplification (WGA) and Rolling Circle Amplification (RCA)

WGA and RCA are examples of Multiple Displacement Amplification (MDA). MDA utilizes the strand-displacement activity of DNA polymerases such as phi29 DNA Polymerase, engineered phi29-XT DNA Polymerase, or *Bst* DNA Polymerase for isothermal amplification of an entire genome or circular templates.

As a processive polymerase that can extend for tens of thousands of nucleotides in a single binding event, as well as increased proofreading activity, phi29-XT DNA Polymerase is ideal for accurate amplification of long or circular templates.


RECOMMENDED PRODUCTS

phi29-XT WGA Kit
(NEB #E1604)

phi29-XT RCA Kit
(NEB #E1603)


T7 Endonuclease I
(NEB #M0302)

Comparison of phi29-XT and phi29 DNA Polymerase yields



Triplicate WGA reactions were performed with 100 pg human genomic DNA input at the indicated temperatures for 1.5 hours. The reactions were then heat-inactivated at 65°C for 10 minutes. All reactions contained 1 mM dNTPs and 50 µM Exonuclease-Resistant Random Primers. Wild-type phi29 reactions were carried out with 10 units of phi29 DNA Polymerase (NEB #M0269) in 1X phi29 DNA Polymerase Reaction Buffer and 0.1 mg/mL Recombinant Albumin, and phi29-XT reactions were carried out with 1X phi29-XT DNA Polymerase for WGA and 1X phi29-XT Reaction Buffer for WGA. Reaction yields (dots) were quantified using Quant-iT[®] PicoGreen[®] dsDNA Reagent and were averaged (lines) to determine the yield at each reaction temperature. While the working reaction temperature of wild-type phi29 DNA Polymerase is typically between 30°C and 38°C, phi29-XT DNA Polymerase generates robust product yields around 42°C. The optimal temperature range for wild-type phi29 increased by approximately three degrees for WGA using human genomic DNA when compared to RCA using pUC19 plasmid as input.

Overview of the phi29-XT WGA Kit



The phi29-XT WGA Kit (NEB #E1604) is a fast, easy-to-use, and highly versatile kit containing all the required components for whole genome amplification (WGA) using a random primer mix. The kit delivers high yields of DNA products from a variety of starting materials including purified genomic DNA, cells, or microbiomes. This kit is ideal for various DNA applications such as single cell whole genome sequencing (WGS), genomic DNA enrichment, and environmental DNA (eDNA) enrichment and sequencing.

Strand Displacement Amplification (SDA)

SDA relies on a strand-displacing DNA polymerase, typically *Bst* DNA Polymerase, Large Fragment (NEB #M0275) or Klenow Fragment (3'-5' exo-) (NEB #M0212) to initiate amplification at nicks created by a strand-limited restriction endonuclease or nicking enzyme (e.g. WarmStart Nt.BstNBI, NEB #R0725) at a site contained in a primer. The nicking site is regenerated with each polymerase displacement step, resulting in exponential amplification.

RECOMMENDED PRODUCTS

WarmStart Nt.BstNBI (NEB #R0725)


***Bst* DNA Polymerase, Full Length** (NEB #M0328)

***Bst* DNA Polymerase, Large Fragment**
(NEB #M0275)

***Bst* 2.0 DNA Polymerase** (NEB #M0537)

WarmStart Afu Uracil-DNA Glycosylase (UDG)
(NEB #M1282)

Overview of SDA




* Target amplification, shown above for SDAF, will also occur simultaneously with SDAR.

For more information on these and other isothermal amplification methods and products, please visit www.neb.com/isoamp

Recombinase Polymerase Amplification (RPA)

Recombinase Polymerase Amplification (RPA) uses a recombinase enzyme to help primers invade double-stranded DNA. T4 UvsX, UvsY, and a single stranded binding protein T4 Gene 32 protein form D-loop recombination structures that initiate amplification by a strand-displacing DNA polymerase. RPA is typically performed at ~37 °C and, unlike other methods, can produce discrete amplicons up to 1 kb.

Overview of RPA



RPA utilizes a recombinase-primer complex, a strand-displacing polymerase, and single-stranded DNA binding proteins to facilitate amplification of discrete DNA products up to 1 kb (primer-to-primer distance).

RECOMMENDED PRODUCTS

T4 UvsX Recombinase (Glycerol-free)*
(NEB #M3081)


T4 UvsY Protein (Glycerol-free)* (NEB #M3082)

Bsu DNA Polymerase, Large Fragment
(NEB #M0330)

T4 Gene 32 Protein (NEB #M0300)

* Please contact the Customized Solutions Team at www.neb.com/customizedsolutions, to inquire about these products.

Resources to support your isothermal amplification work



Download our ebook, featuring:

- Comparison of PCR and isothermal amplification technologies
- Examination of assay design optimization parameters
- Considerations for assay lyophilization
- Testimonial from Anne Wyllie, Ph.D.,
Research Scientist in Epidemiology of Microbial Disease, Yale School of Public Health



Register here to receive
your copy of the eBook



NEB LAMP Primer Design

To begin, use our free online tool to design primers
for any of these methods at lamp.neb.com

What products are recommended for molecular diagnostics?

NEB offers a selection of reverse transcriptases (RTs), DNA polymerases, and master mixes for customers to use in the development of rapid and sensitive molecular diagnostics. To support the needs of this market, products are available in fully lyophilized, high concentration, or glycerol-free/lyo-compatible formats. All formats are available for customization.

WarmStart RTx is ideal for LAMP and isothermal amplification workflows and is a core component of a lyophilized LAMP/RT-LAMP mix (NEB #L4401). Luna® RT supports RT-qPCR workflows and is available in a lyophilized format (NEB #L4001). Customers have utilized these enzymes in numerous diagnostic assays, including for COVID-19.

NEB Lyophilization Sciences® is equipped to develop and manufacture lyophilized molecular biology reagents for the life sciences, including research, applied and the molecular diagnostics sectors. The NEB Lyophilization Sciences Team are experts in the design, development and manufacturing of innovative solutions for ambient-stored products.



Products can be lyophilized in multiple formats.

Contact us at www.neb.com/customizedsolutions
to find out how we can support your development,
or visit www.neb.com/mdx to learn more.

Choose from our selection of products for your isothermal application.

PRODUCT	NEB #	SIZE
WarmStart Multi-Purpose LAMP/RT-LAMP 2X Master Mix (with UDG)	M1708S/L	100/500 reactions
WarmStart Fluorescent LAMP/RT-LAMP Kit (with UDG)	E1708S/L	100/500 reactions
WarmStart LAMP Kit (DNA & RNA)	E1700S/L	100/500 reactions
LAMP Fluorescent Dye	B1700S	0.25 ml
WarmStart Colorimetric LAMP 2X Master Mix (DNA & RNA)	M1800S/L	100/500 reactions
WarmStart Colorimetric LAMP 2X Master Mix with UDG	M1804S/L	100/500 reactions
SARS-CoV-2 Rapid Colorimetric LAMP Assay Kit	E2019S	96 reactions
Bst-XT WarmStart DNA Polymerase	M9204S/L	1,600/8,000 units
Bst 3.0 DNA Polymerase	M0374S/L/M	1,600/8,000/8,000 units
Bst 2.0 WarmStart DNA Polymerase	M0538S/L/M	1,600/8,000/8,000 units
Bst 2.0 DNA Polymerase	M0537S/L/M	1,600/8,000/8,000 units
Bst DNA Polymerase, Large Fragment	M0275S/L/M	1,600/8,000/8,000 units
Bst DNA Polymease, Full Length	M0328S	500 units
phi29-XT RCA Kit	E1603S/L	100/500 reactions
phi29-XT WGA Kit	E1604S/L	25/100 reactions
WarmStart Afu Uracil-DNA Glycosylase (UDG)	M1282S	200 units
WarmStart Nt.BstNBI	R0725S	1,000 units
WarmStart RTx Reverse Transcriptase	M0380S/L	50/250 reactions
Nt.BstNBI	R0607S/L	1,000/5,000 units
COMPANION PRODUCTS		
Tte UvrD Helicase	M1202S	0.5 µg
Deoxynucleotide (dNTP) Solution Mix	N0447S/L	8/40 µmol of each
Deoxynucleotide (dNTP) Solution Set	N0446S	25 µmol of each

Products and content are covered by one or more patents, trademarks and/or copyrights owned or controlled by New England Biolabs, Inc. (NEB). The use of trademark symbols does not necessarily indicate that the name is trademarked in the country where it is being read; it indicates where the content was originally developed. All other trademarks are the property of their respective owners. The use of this product may require the buyer to obtain additional third-party intellectual property rights for certain applications. For more information, please email busdev@neb.com.

Your purchase, acceptance, and/or payment of and for NEB's products is pursuant to NEB's Terms of Sale at www.neb.com/support/terms-of-sale. NEB does not agree to and is not bound by any other terms or conditions, unless those terms and conditions have been expressly agreed to in writing by a duly authorized officer of NEB.

Licensed under U.S. Patents Nos. U.S. 6,410,278; 6,974,670; 7,494,790; 8,017,357; 7,374,913; 7,851,186; and 7,846,695 including all foreign counterparts thereof. The purchase of this product conveys to the purchaser the limited, non-transferable right to use the purchased product to perform loop-mediated isothermal amplification ("LAMP") for Research Use Only. The purchase of this product further conveys to the purchaser the limited, non-transferable right to use the purchased product to perform reverse transcription loop-mediated isothermal amplification ("RT-LAMP") for Research Use Only. LAMP is a patented technology belonging to Eiken Chemical Co., Ltd. and any use other than research may require a license from Eiken Chemical Co., Ltd. Notice to Purchaser: Nucleic-acid based aptamers for use with thermophilic DNA polymerases are licensed exclusively by New England Biolabs, Inc. from Somalogic, Inc. New England Biolabs, Inc. gives the Buyer/User a non-exclusive license to use the aptamer-based RT-LAMP Master Mix for Research Purposes Only. Commercial use of the aptamer-based RT-LAMP Master Mix requires a license from New England Biolabs, Inc. Please contact busdev@neb.com for more information.

The purchase of NEB Bst products conveys to the purchaser the limited, nontransferable right to use the purchased products to perform loop-mediated isothermal amplification ("LAMP") for research use only. LAMP is a patented technology belonging to Eiken Chemical Co., Ltd. and any use other than research may require a license from Eiken Chemical Co., Ltd.

The purchase of NEB RTx products conveys to the purchaser the limited, non-transferable right to use the purchased products to perform reverse transcription loop-mediated isothermal amplification ("RT-LAMP") for research use only. LAMP is a patented technology belonging to Eiken Chemical Co., Ltd. and any use other than research may require a license from Eiken Chemical Co., Ltd. This product is intended for research purposes only. This product is not intended to be used for therapeutic or diagnostic purposes in humans or animals.

B CORPORATION® is a registered trademark of B Lab Company
Quant-IT® is a registered trademark of Life Technologies Corporation
PicoGreen® is a registered trademark of Molecular Probes, Inc.

© Copyright 2025, New England Biolabs, Inc.; all rights reserved.



Did you know that many of these products can be purchased in large volumes and custom formats?
Learn more at www.neb.com/customizedsolutions

x/E/yNEB123_V5.0_0725

GERMANY & AUSTRIA

New England Biolabs GmbH
Brüningstr. 50, Geb B852
65926 Frankfurt/Main, Germany

Tel: +49/(0)69/305-23140
Fax: +49/(0)69/305-23149
Free Call: 0800/246 5227 (Germany)
Free Call: 00800/246 52277 (Austria)
info.de@neb.com

www.neb-online.de

HEADQUARTERS: USA

New England Biolabs, Inc.

Telephone: (978) 927-5054
Toll Free (USA Orders): 1-800-632-5227
Toll Free (USA Tech): 1-800-632-7799
Fax: (978) 921-1350
info@neb.com

www.neb.com

FRANCE

New England Biolabs France SAS
Genopole Campus 1, Bâtiment 6
5 rue Henri Desbruères
91030 Evry cedex, France

Tel.: 0800 100 632 (Customer Service)
Tel.: 0800 100 633 (Technical Service)
info.fr@neb.com

www.neb-online.fr

For a list of NEB subsidiaries and distributors, please go to:

www.neb.com/international-support



Print product with financial
climate contribution
ClimatePartner.com/13845-2508-1009

